

- Write your **name** on every sheet, and on the first sheet your **student number, group** (1: Aldo and Francesco, 2: Maarten) and the total **number of sheets** handed in.
- Use a **separate sheet** for each exercise!
- You may use the lecture notes, the extra notes and personal notes, but no worked exercises.
- Do not just give answers, but also justify them with complete arguments. If you use results from the lecture notes, always **mention this**, and show that their hypotheses are fulfilled in the situation at hand.
- **N.B.** If you fail to solve an item within an exercise, **do continue**; you may then use the information stated earlier.
- The weights by which exercises and their items count are indicated in the margin. The highest possible total score is 44. The grade will be obtained from your total score  $T$  by rounding off  $\min(T/4, 10)$  to a half integer above 6 or an integer below 6.5.
- You are free to write the solutions either in English, or in Dutch.

*Succes !*

13 pt total **Exercise 1.** For  $\mathbb{R}$  we consider the collection  $\mathcal{B}$  of all subsets of the form

$$[p, q) := \{x \in \mathbb{R} \mid p \leq x < q\}, \quad \text{with } p, q \in \mathbb{Q}, p < q.$$

- 3 pt (a) Show that  $\mathcal{B}$  is not a topology, but it is a topology basis. Denote by  $\mathcal{T}$  the topology generated by  $\mathcal{B}$ .
- 2 pt (b) Show that  $\mathcal{T}$  contains the Euclidean topology.
- 2 pt (c) Is  $(\mathbb{R}, \mathcal{T})$  first countable, second countable? Is it Hausdorff?
- 3 pt (d) Show that  $[0, 1)$  is open and closed in  $(\mathbb{R}, \mathcal{T})$ . Is  $(\mathbb{R}, \mathcal{T})$  connected?
- 3 pt (e) Show that  $[0, 1)$  and  $[0, 1]$  are not compact in  $(\mathbb{R}, \mathcal{T})$ .

9 pt total **Exercise 2.** Let  $X := [0, 1] \times [-2, 2]$  and the subset  $B := [0, 1] \times [-1, 1] \subset X$  both be equipped with the topology induced by the Euclidean topology on  $\mathbb{R}^2$ .

We equip  $X$  with the equivalence relation  $R$  whose equivalence classes are given by  $\{(s, t)\}$  for  $0 < s < 1$  and  $-2 < t < 2$ ,  $\{(s, -2), (s, 2)\}$  for  $0 < s < 1$ ;  $\{(0, t), (1, -t)\}$  for  $-2 < t < 2$  and, finally,  $\{(0, \pm 2), (1, \pm 2)\}$ . Accordingly,  $X/R$  equipped with the quotient topology, is the Klein bottle (which is known to be a Hausdorff space). The associated quotient map is denoted by  $\pi : X \rightarrow X/R$ .

- 3 pt (a) The restriction of  $R$  to  $B$  is the relation  $R_B$  defined by  $b_1 R_B b_2 \iff \pi(b_1) = \pi(b_2)$ , for  $b_1, b_2 \in B$ . Show that  $R_B$  is an equivalence relation on  $B$  and explicitly determine the associated equivalence classes in  $B$ .
- 1 pt (b) The quotient  $B/R_B$  is equipped with the quotient topology. To which well known space is this quotient homeomorphic? (You need not justify your answer.)

Let  $\pi_B : B \rightarrow B/R_B$  be the associated quotient map.

- 2 pt (c) Show that there exists a unique map  $f : B/R_B \rightarrow X/R$  such that for all  $b \in B$  we have  $f(\pi_B(b)) = \pi(b)$ .
- 3 pt (d) Prove that the map  $f$  is an embedding.

11 pt total **Exercise 3.** Let  $M$  be a topological space, and assume that  $\gamma \mapsto \varphi_\gamma$  is an action of the group  $\mathbb{Z}_2 = \{-1, 1\}$  on  $M$  by homeomorphisms. Let  $M/\mathbb{Z}_2$  be the associated quotient (equipped with the quotient topology), and  $\pi : M \rightarrow M/\mathbb{Z}_2$  the quotient map.

- 2 pt (a) Given a subset  $V \subset M$ , show that  $\pi^{-1}(\pi(V)) = V \cup \varphi_{-1}(V)$ .
- 2 pt (b) Show that for  $V \subset M$  open, the set  $\pi(V)$  is open in  $M/\mathbb{Z}_2$ .
- 3 pt (c) Let  $\{U_i\}_{i \in I}$  be an open cover of  $M$ . For every  $m \in M$  let  $i_m, j_m \in I$  be indices such that  $m \in U_{i_m}$  and  $\varphi(m) \in U_{j_m}$ . Show that there exists an open neighborhood  $V_m$  of  $m$  such that  $V_m \subset U_{i_m}$  and  $\varphi(V_m) \subset U_{j_m}$ .
- 4 pt (d) Show that  $M$  is compact if and only if  $\pi(M)$  is compact. Hint: for one of the implications consider the collection  $\{\pi(V_m)\}$  resulting from (c).

11 pt total **Exercise 4.** For  $M$  a locally compact Hausdorff space we denote by  $C_c(M)$  the real linear space of continuous functions  $M \rightarrow \mathbb{R}$  with compact support. If  $U$  is an open subset of  $M$ , we put  $C_c(U) := \{f \in C_c(M) \mid \text{supp} f \subset U\}$ .

- 2 pt (a) If  $U$  is open in  $M$ ,  $f \in C_c(M)$ ,  $\psi \in C(M)$  and  $\text{supp} \psi \subset U$ , show that  $\psi f \in C_c(U)$ .

By a positive integral on an open subset  $U$  of  $M$  we mean a linear map  $I : C_c(U) \rightarrow \mathbb{R}$  such that for all  $f \in C_c(U)$  we have:

$$f \geq 0 \Rightarrow I(f) \geq 0.$$

A positive integral  $I$  on  $U$  is said to be strictly positive if for all  $f \in C_c(U)$  we have

$$f \geq 0, I(f) = 0 \Rightarrow f = 0.$$

- 1 pt (b) Prove the following result. If  $I$  is a positive integral on an open subset  $U$  of  $M$  and  $\psi : M \rightarrow \mathbb{R}$  is continuous function with  $\psi \geq 0$  and  $\text{supp} \psi \subset U$  then  $I_\psi : f \mapsto I(\psi f)$  is a positive integral on  $M$ .
- 5 pt (c) Assume that  $M$  is second countable, and that for every point  $m \in M$  there exists an open neighborhood  $U_m \ni m$  and a strictly positive integral  $I_m : C_c(U_m) \rightarrow \mathbb{R}$  on  $U_m$ . Show that there exists a strictly positive integral on  $M$ .

In the next item you may use without proof that the map  $J : C_c((0, 1)^n) \rightarrow \mathbb{R}$  defined by the  $n$ -fold repeated Riemann integral  $J(f) = \int_0^1 \cdots \int_0^1 f(x) dx_1 \cdots dx_n$ , is a strictly positive integral on the open subset  $(0, 1)^n$  of  $\mathbb{R}^n$ .

- 3 pt (d) If  $M$  is a topological manifold of dimension  $n \geq 1$ , show that there exists a strictly positive integral on  $M$ .